



Chemistry Notes for class 12 Chapter 13

Amines

Amines constitute an important class of organic compounds derived by replacing one or more hydrogen atoms of NH_3 molecule by alkyl/aryl group(s).




In the IUPAC system, the amines are regarded as alkanamines, e.g.,




Structure


The nitrogen atom in amine is sp^3 -hybridised. The three hybrid orbitals are involved in bond formation and one hybrid atomic orbital contains the lone pair of electrons, giving the pyramidal geometry of amines.




In arylamines, $-\text{NH}_2$ group is directly attached to the benzene ring.






Methods of Preparation of Amines

(i) Reduction of nitro compounds

Reduction can take place by Sn/HCl, Ni/H₂, Zn/NaOH, Pd/H₂.



The reduction of nitroalkane or nitrobenzene in neutral medium gives hydroxyl amines.



(ii) Ammonolysis of alkyl halides

Ammonolysis has the disadvantage of yielding a mixture of primary, secondary and tertiary amines and also a quaternary ammonium salt.



However, primary amine is obtained as a major product by taking large excess of NH₃.

Order of reactivity of halides with amines is RI > RBr > RCl.


Aromatic amines could not be prepared since aryl halides are much less reactive towards nucleophilic substitution reactions.

(iii) Reduction of nitriles or cyanides**(iv) Schmidt reaction**

It is a modification of Curtius degradation.

(v) Reduction of amides**(vi) Gabriel's phthalimide reaction**

It only produces 1° amines. This method is not suitable for 1° arylamine because aryl halide does not give nucleophilic substitution reaction.

(viii) Hofmann bromamide degradation reaction

In Hofmann degradation reaction, the amine formed has one carbon less than the parent amide. To obtain primary amine with same number of carbon atoms from primary amide, reduction is done with $\text{LiAlH}_4/\text{ether}$.

Physical Properties of Amines

1. The lower aliphatic amines are gases with fishy smell.
2. Primary amines with three or more carbon atoms are liquid and higher members are all solids.
3. Lower aliphatic amines are water soluble because they can form hydrogen bonds with water molecules, however the solubility decreases with increase in hydrophobic alkyl group.
4. Boiling points order primary > secondary > tertiary

5. Tertiary amines does not have intermolecular association due to the absence of hydrogen atom available for hydrogen bond formation.

Basic Strength of Amines

Amines act as Lewis bases due to the presence of lone pair of electrons on the nitrogen atom.

More the K_b (dissociation constant of base), higher is the basicity of amines.

Lesser the pK_b , higher is the basicity of amines.

Aliphatic amines (CH_3NH_2) are stronger bases than NH_3 due to the electron releasing +/ effect of the alkyl group.

Among aliphatic methyl amines, the order of basic strength in aqueous solution is as follows




Aromatic amines are weaker bases than aliphatic amines and NH_3 , due to the fact that the electron pair on the nitrogen atom is involved in resonance with the π -electron pairs of the ring.

Electron releasing groups (e.g., $-\text{CH}_3$, $-\text{OCH}_3$, $-\text{NH}_2$ etc.) increase the basic strength of aromatic amines while electron withdrawing groups (like $-\text{NO}_2$, $-\text{X}$, $-\text{CN}$ etc.) tend to decrease the same.

o-substituted aromatic amines are usually weaker bases than aniline irrespective of the nature of substituent whether electron releasing or electron withdrawing. This is called ortho effect and is probably due to steric and electronic factors.

chemical Properties of Amines

(i) **Alkylation** All the three types of amines react with alkyl halides to form quaternary ammonium salt as the final product provided alkyl halide is present in excess.




Aromatic amines also undergo alkylation as given below.


But secondary and tertiary amines react with nitrous acid in different manner.

Methyl amine give dimethyl ether with HNO_2 .

(vi) Reaction with aryl sulphonyl chloride [Hinsberg reagent] The reaction of benzenesulphonyl chloride with primary amine yield N-ethyl benzenesulphonyl amide.




The reaction of benzenesulphonyl chloride with secondary amine yields N,N-diethyl benzene sulphonamide.




Tertiary amines does not react with benzenesulphonyl chloride.


(vii) Reaction with aldehydes Schiff base is obtained.




(viii) Electrophilic substitution reactions Aniline is ortho and para directing towards electrophilic substitution reaction due to high electron density at ortho and para-positions.




To prepare monosubstituted derivative, activating effect of $-NH_2$ group must be controlled. It can be done by protecting the $-NH_2$ group by acetylation with acetic anhydride.




(b) **Nitration Direct** nitration of aniline is not possible as it is susceptible to oxidation, thus amino group is first protected by acetylation.



In strongly acidic medium, aniline is protonated as anilinium ion which is meta directing so it gives meta product also.



(c) **Sulphonation** On sulphonation, aniline gives sulphanilic acid, as the major product.



(d) Aniline does not undergo Friedel-Crafts reaction due to salt formation with aluminium chloride, the Lewis acid, which is used as a catalyst. Due to this, nitrogen of aniline acquires positive charge and hence, behave like a strong deactivating group for further chemical reaction.

(ix) **Oxidation** Use of different oxidising agents gives different products.

e.g.,

Oxidising agent	Product
Acidified KMnO_4 (or $\text{Na}_2\text{C}_2\text{O}_7 + \text{CuSO}_4 + \text{dil acid}$)	Aniline black (a dye)
Chromic acid ($\text{Na}_2\text{C}_2\text{O}_7 + \text{Conc H}_2\text{SO}_4$)	p-benzoquinone
Caro's acid (H_2SO_5)	nitrobenzene and nitrosobenzene
Conc. nitric acid	decomposes

Separation of Mixture of Amines (1° , 2° and 3°)

(a) Fractional distillation This method is based on the boiling points of amines and is used satisfactorily in industry.


(b) Hofmann's method Diethyloxalate is called Hofmann's reagent with which mixture of amines is treated.

- 1° amine forms solid dialkyl oxamide (CONHR)₂
- 2° amine forms liquid dialkyl oxamic ester (CONR₂-COOC₂H₅)
- 3° amines do not react

(c) Hinsberg's method see chemical reactions.

Benzene Diazonium Chloride (C₆H₅N₂⁺;Cl⁻)

Preparation (Diazotisation reaction)



The excess acid in diazotisation reaction is necessary to maintain proper acidic medium for the reaction and to prevent combination of diazonium salt formed with the undiazotised amine.


Diazonium salts are prepared and used in aqueous solutions because in solid state, they explode.

Properties

It is a colourless crystalline solid, soluble in water. It has tendency to explode when dry.


Stability of Arenediazonium salts


It is relatively more stable than the alkyldiazonium salt. The arenediazonium ion is resonance stabilised as is indicated by the following resonating structures:



Various resonating structures of arenediazonium ion

Chemical Reactions






Alkyl Cyanides'


These compound have formula RCN. These are the derivatives of RCN.

According to IUPAC system, cyanides are named as 'alkane nitrile', e.g.,




Methods of Preparation

(i) From alkyl halides



(ii) From acid amides




Physical properties


1. These are neutral compound with pleasant odour, similar to bitter almonds.
2. These are soluble in water as well as organic solvents.
3. These are poisonous but less than HCN.

Chemical Properties


(i) Hydrolysis



(ii) Reduction



(iii) Reaction with Grignard reagent



Alkyl isocyanides (RNC)


According to IUPAC system, these are named as 'alkane isonitrile'

e.g., CH_3NC methyl isonitrile


C_6H_5NC benzene isonitrile

Methods of Preparation


(a) From alkyl halides



(b) Carbylamine reaction



(c) From N-alkyl formamide




Physical Properties


1. These are colourless unpleasant smelling liquids.
2. These are soluble in organic solvents but insoluble in water.

Chemical Properties

(i) Hydrolysis




(ii) Reduction



(iii) **Addition reaction** Due to the presence of unshared pair of electrons on C atom, alkyl isocyanides give addition reaction.



(iv) **Isomerisation** At 250°C, it isomerises to nitrile.



Nitro Compounds

These are obtained by replacing one H of hydrocarbon by $-\text{NO}_2$ group.


These are named according to IUPAC system as 'nitro alkane'.

Methods of Preparation

(i) From alkyl halides



(ii) **Nitration** Nitrating mixture is conc HNO_3 + conc H_2SO_4 .



Physical Properties


1. These are colourless pleasant smelling liquids.
2. Their boiling point are much higher than isomeric alkyl nitriles.
3. These are less soluble in water but readily soluble in organic solvents.

Chemical Properties


(i) **Reduction** With Sn/HCl or catalytic hydrogenation, nitroalkanes are reduced to amines.



If neutral reducing agent like Zn dust + NH_4Cl is used, hydroxylamines are obtained as major product.



In the presence of $(NH_4)_2S$ or Na_2S , selective reduction takes place.




Nitrobenzene gives different products with different reagents and in different mediums.


Medium	Reagent	Product
Acid	Sn/HCl	aniline
Neutral	Zn/NH_4Cl	N-phenyl hydroxylamine
Alkaline	$Na_3AsO_3/NaOH$	azoxybenzene ($C_6H_5N=O-C_6H_5$)
	$Zn/NaOH, CH_3OH$	azobenzene
Metallic hydride	$Zn/NaOH, C_2H_5OH$	hydrazobenzene
Electrolytic	$LiAlH_4$	aniline
	dil H_2SO_4	p-aminophenol

(ii) Action of HNO_2

1° nitroalkane gives nitrolic acid which gives red colour with $NaOH$.




2° nitroalkanes give pseudonitrol with HNO_2 .




3° nitroalkanes does not react with HNO_2

(iii) **Nef carbonyl** synthesis Na or K salt of 1° or 2° nitroalkanes give carbonyl compounds on acidification with 50% H_2SO_4 at room temperature. This reaction is called Nef carbonyl synthesis.



(iv) **Electrophilic** substitution On nitration, nitrobenzene gives m-dinitrobenzene (as $-\text{NO}_2$ is a m-directing group and strongly deactivating).



It does not give Friedel-Craft's alkylation.

(v) **Nucleophilic** substitution reaction $-\text{NO}_2$ group activates the ring towards nucleophilic substitution.

